

iF647 Anti-mouse/human Helios Antibody

Catalog Number:	204603, 204604
Size:	25 tests, 100 tests
Target Name:	Helios, IKAROS family zinc finger 2, IKZF2, ZNF1A2, ZNFN1A2
Regulatory Status:	RUO

PRODUCT DETAILS

Clone:	22F6
Application:	Intracellular Flow Cytometry
Reactivity:	Mouse, Human
Format:	iF647
Isotype:	Armenian Hamster IgG
Antibody Type:	Monoclonal
Formulation:	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 0.2% (w/v) BSA
Protein Concentration:	Supplied at a lot-specific concentration.
Storage&Handling:	The antibody solution should be stored undiluted between 2°C and 8°C, and protected from prolonged exposure to light. Do not freeze.
Recommended Usage:	For flow cytometric staining, it is recommended to use less than 0.2 µg of this reagent per 0.5-1.0 million cells in a 100 µL volume. Optimal reagent performance should be determined by titration for each specific application. iF647 has an excitation max at 656 nm and an emission max at 670 nm.
Excitation Laser:	Red Laser (633 nm)
Isotype Control:	300511
Isotype Controls:	300511

BACKGROUND INFORMATION


Helios, encoded by the IKZF2 gene, is a transcription factor belonging to the Ikaros family of zinc finger proteins. It is predominantly expressed in T lymphocytes, especially regulatory T cells (Tregs), where it contributes to immune homeostasis. Helios is thought to help stabilize Treg identity and suppressive function, although its exact role in distinguishing thymic versus peripherally induced Tregs remains an area of active investigation.

Structurally, Helios contains multiple C2H2 zinc finger domains that mediate DNA binding and protein-protein interactions. The N-terminal zinc fingers are responsible for sequence-specific binding to target gene promoters, while the C-terminal domains facilitate dimerization with other Ikaros family members. As a nuclear protein, Helios does not interact with extracellular ligands; instead, it regulates transcriptional programs controlling T-cell activation and differentiation.

In disease, dysregulation of Helios has been associated with autoimmune disorders, where impaired Treg function may contribute to loss of immune tolerance. Altered Helios expression is also observed in certain T-cell leukemias and lymphomas, suggesting a role in malignant transformation or persistence.

Therapeutically, Helios is not directly targeted but is an important biomarker and functional regulator in immunotherapy. Modulating Helios expression or Helios-positive Tregs may enhance anti-tumor immunity, particularly in cancer settings where Tregs suppress effective immune responses. Ongoing research aims to exploit this pathway while preserving immune balance.

PRODUCT DATA



Human peripheral blood lymphocytes were stained with iF488 anti-Human CD4 clone 004AB and iF647Anti-Mouse/Human Helios clone 22F6..

This product is supplied subject to the terms and conditions at www.innocyto.com/web/terms.php and may only be used as provided in the stated terms. Products are for Research Use Only.