

## Biotin Human BCMA (CD269) Protein (C-Fc-Avi)

<b>Catalog Number:</b>	802203, 802204
<b>Size:</b>	25 ug, 100 ug
<b>Target Name:</b>	TNFRSF17, CD269, BCM, BCMA
<b>Regulatory Status:</b>	RUO

### PRODUCT DETAILS

---

<b>Application:</b>	ELISA, BLI
<b>Format:</b>	Liquid, Biotinylated
<b>Expression Host:</b>	HEK293
<b>Species:</b>	Human
<b>Sources:</b>	Human BCMA protein (NP_001183.2) (Met1-Ala54) with C-terminus Human IgG1 Fc-Avi tag is expressed in HEK293 cells. This protein was site-specifically labeled with Biotin by BirA ligase.
<b>Accession Number:</b>	Q02223
<b>Molecular Weight:</b>	The protein has a predicted molecular weight of 34.3kDa. Under DTT-reducing conditions, it migrates at approximately 35-45 kDa on SDS-PAGE.
<b>Affinity Tag:</b>	C-Fc-Avi
<b>Purity:</b>	>95% based on SDS-PAGE under reducing condition
<b>Formulation:</b>	1xPBS buffer, pH7.4, 0.22 µm filtered
<b>Endotoxin level:</b>	Not tested
<b>Protein Concentration:</b>	25µg size is bottled at 0.2mg/mL concentration. 100 µg size is supplied at a lot-specific concentration.
<b>Storage and Handling:</b>	Briefly centrifuge the vial upon receipt. An unopened vial can be stored at 4°C for up to 2 weeks, or at -20°C or below for up to six months. The protein may be further diluted to 0.1 mg/mL using 0.22 µm-filtered PBS buffer (pH 7.4). For long-term storage, the diluted stock solution should be aliquoted and stored at ≤ -70°C to minimize freeze-thaw cycles. If additional dilution is required, carrier proteins such as FBS or BSA should be added to maintain protein stability.


### BACKGROUND INFORMATION

---

CD269, also known as B-cell maturation antigen (BCMA), is a 27 kD single-pass transmembrane protein that contains one TNFR-Cys repeat in its extracellular domain. This protein is a critical factor for the long-term survival and maintenance of plasma cells and plays an essential role in B-cell maturation. CD269 is expressed by plasmablasts, plasma cells, and germinal center B cells, and it is primarily involved in regulating humoral immunity. The ligands for CD269 include BAFF (B-cell activating factor) and APRIL (A proliferation-inducing ligand). These ligands bind to CD269, initiating a signaling cascade that supports B-cell survival and plasma cell function. Additionally, the cytoplasmic domain of CD269 interacts with several members of the TRAF (tumor necrosis factor

receptor-associated factor) family, which are important for transducing survival and proliferation signals within the cell. Due to its essential role in immune response regulation, CD269 is a key player in the development and functioning of plasma cells.

## PRODUCT DATA



Human BCMA (C-Fc-Avi) was biotinylated in vitro using BirA ligase. SDS-PAGE analysis under reducing (P+) and non-reducing (P-) conditions shows the protein has a purity greater than 95%. A gel shift assay using co-incubation with streptavidin indicates that the biotinylation efficiency of the BCMA protein exceeds 95%.

This product is supplied subject to the terms and conditions at [www.innocyto.com/web/terms.php](http://www.innocyto.com/web/terms.php) and may only be used as provided in the stated terms. Products are for Research Use Only.