

Human IL13RA2 (CD213A2) Protein (C-Fc-Avi)

Catalog Number:	804801, 804802
Size:	25 ug, 100 ug
Target Name:	IL13RA2, CD213A2, IL-13R, IL13BP
Regulatory Status:	RUO

PRODUCT DETAILS


Application:	ELISA, BLI
Format:	Liquid, Purified
Expression Host:	CHO
Species:	Human
Sources:	Recombinant Human IL13RA2 protein (Cys22-Leu342) with C-terminus Fc-Avi tag is expressed in CHO cells.
Accession Number:	Q14627
Molecular Weight:	The protein has a predicted molecular weight of 65.7 kDa. Under DTT-reducing conditions, it migrates at approximately 65-80 kDa on SDS-PAGE.
Affinity Tag:	C-Fc-Avi
Purity:	>95% based on SDS-PAGE under reducing condition
Formulation:	1xPBS buffer, pH7.4, 0.22 µm filtered
Endotoxin level:	Not tested
Protein Concentration:	25µg size is bottled at 0.2mg/mL concentration. 100 µg size is supplied at a lot-specific concentration.
Storage and Handling:	Briefly centrifuge the vial upon receipt. An unopened vial can be stored at 4°C for up to 2 weeks, or at -20°C or below for up to six months. The protein may be further diluted to 0.1 mg/mL using 0.22 µm-filtered PBS buffer (pH 7.4). For long-term storage, the diluted stock solution should be aliquoted and stored at ≤ -70°C to minimize freeze-thaw cycles. If additional dilution is required, carrier proteins such as FBS or BSA should be added to maintain protein stability.

BACKGROUND INFORMATION

CD213A2, also known as IL13Rα2, is a type I transmembrane protein belonging to the hematopoietin receptor family. It binds interleukin-13 (IL-13) with high affinity but lacks a functional cytoplasmic signaling domain, suggesting that it primarily acts as a decoy receptor, antagonizing IL-13 signaling mediated by the IL-13Rα1/IL-4Rα complex. IL13Rα2 is expressed in fibroblasts, smooth muscle cells, keratinocytes, and activated B cells, though its surface expression is tightly regulated and much of it resides intracellularly or in soluble form. In addition to inhibiting IL-13 activity, IL13Rα2 has been reported to suppress IL-4 signaling via physical interaction with IL-4Rα, while paradoxically promoting TGF-β production and fibrosis. Importantly, IL13Rα2 is highly and

specifically overexpressed in certain cancers, such as glioblastoma multiforme and high-grade astrocytomas, making it a promising target for tumor-specific immunotherapies and viral vector-mediated gene delivery approaches.

PRODUCT DATA



Human IL13RA2 Protein (C-Fc-Avi) on SDS-PAGE under reducing condition (P+) and non-reducing condition (P-). The gel was stained for 1 hour with BlinkBlue (catalog 700102). The purity of this protein appears to be greater than 95% based on reducing conditions.

This product is supplied subject to the terms and conditions at www.innocyto.com/web/terms.php and may only be used as provided in the stated terms. Products are for Research Use Only.