

## Human CD19 Protein (C-Fc)

<b>Catalog Number:</b>	807301, 807302
<b>Size:</b>	25 ug, 100 ug
<b>Target Name:</b>	CD19, B4, CVID3
<b>Regulatory Status:</b>	RUO

### PRODUCT DETAILS

---

<b>Application:</b>	ELISA, BLI
<b>Format:</b>	Liquid, Purified
<b>Expression Host:</b>	CHO
<b>Species:</b>	Human
<b>Sources:</b>	Recombinant Human CD19 Protein (Glu21-Lys291) with C-terminus Fc-tag is expressed in CHO cell.
<b>Accession Number:</b>	P15391
<b>Molecular Weight:</b>	The protein has a predicted molecular weight of 56.2 kDa. Under DTT-reducing conditions, it migrates at approximately 70-80 kDa on SDS-PAGE.
<b>Affinity Tag:</b>	C-Fc
<b>Purity:</b>	>95% based on SDS-PAGE under reducing condition
<b>Formulation:</b>	1xPBS buffer, pH7.4, 0.22 µm filtered
<b>Endotoxin level:</b>	Not tested
<b>Protein Concentration:</b>	25µg size is bottled at 0.2mg/mL concentration. 100 µg size is supplied at a lot-specific concentration.
<b>Storage and Handling:</b>	Briefly centrifuge the vial upon receipt. An unopened vial can be stored at 4°C for up to 2 weeks, or at -20°C or below for up to six months. The protein may be further diluted to 0.1 mg/mL using 0.22 µm-filtered PBS buffer (pH 7.4). For long-term storage, the diluted stock solution should be aliquoted and stored at ≤ -70°C to minimize freeze-thaw cycles. If additional dilution is required, carrier proteins such as FBS or BSA should be added to maintain protein stability.

### BACKGROUND INFORMATION


---

Cluster of Differentiation 19 (CD19) is a 95 kDa type I transmembrane glycoprotein and a member of the immunoglobulin superfamily, expressed on B cells from the early pro-B stage through to activated B blasts, but absent on plasma cells. It is also present on follicular dendritic cells. CD19 functions as a crucial co-receptor in B cell signaling by forming a complex with CD21 (CR2) and CD81 (TAPA-1), which enhances the sensitivity of B cells to antigens by lowering the activation threshold. Upon antigen binding, CD19 is phosphorylated, recruits Src-family kinases and PI3K, and internalizes with surface immunoglobulin (sIg), promoting downstream signal transduction necessary for B cell development, activation, and differentiation. Mutations in CD19 are

linked to immunodeficiency syndromes with impaired antibody production, and CD19 is a key target in immunotherapy, particularly CAR-T cell treatment of B-cell malignancies.

## PRODUCT DATA

---



Human CD19 Protein (C-Fc) on SDS-PAGE under reducing condition (P+) and non-reducing condition (P-). The gel was stained for 1 hour with BlinkBlue (catalog 700102). The purity of this protein appears to be greater than 95%.

This product is supplied subject to the terms and conditions at [www.innocyto.com/web/terms.php](http://www.innocyto.com/web/terms.php) and may only be used as provided in the stated terms. Products are for Research Use Only.