

Biotin Human CD4 (C-His-Avi)

Catalog Number:	810703, 810704
Size:	25 ug, 100 ug
Target Name:	CD4, CD4mut, LEU3
Regulatory Status:	RUO

PRODUCT DETAILS


Application:	ELISA, BLI
Format:	Liquid, Biotinylated
Expression Host:	CHO
Species:	Human
Sources:	Recombinant Human CD4 Protein (Lys26-Trp390) with C-terminus His-Avi-tag is expressed in CHO cell. This protein was site-specifically labeled with Biotin by BirA ligase.
Accession Number:	P01730
Molecular Weight:	The protein has a predicted molecular weight of 44.3 kDa. Under DTT-reducing conditions, it migrates at approximately 55-60 kDa on SDS-PAGE.
Affinity Tag:	C-His-Avi
Purity:	>95% based on SDS-PAGE under reducing condition
Formulation:	1xPBS buffer, pH7.4, 0.22 µm filtered
Endotoxin level:	Not tested
Protein Concentration:	25µg size is bottled at 0.2mg/mL concentration. 100 µg size is supplied at a lot-specific concentration.
Storage and Handling:	Briefly centrifuge the vial upon receipt. An unopened vial can be stored at 4°C for up to 2 weeks, or at -20°C or below for up to six months. The protein may be further diluted to 0.1 mg/mL using 0.22 µm-filtered PBS buffer (pH 7.4). For long-term storage, the diluted stock solution should be aliquoted and stored at ≤ -70°C to minimize freeze-thaw cycles. If additional dilution is required, carrier proteins such as FBS or BSA should be added to maintain protein stability.

BACKGROUND INFORMATION

CD4 is a single-pass type I transmembrane glycoprotein belonging to the immunoglobulin superfamily, composed of one Ig-like V-type and three Ig-like C2-type domains. It is expressed on helper T cells, regulatory T cells, monocytes, macrophages, and dendritic cells, functioning as a co-receptor for the T-cell receptor (TCR) by binding MHC class II molecules on antigen-presenting cells to enhance TCR signaling. CD4 also serves as the primary receptor for HIV-1, binding to viral gp120 to mediate viral entry. HIV-1 viral protein U (VpU) binds to the membrane-proximal region of CD4's cytoplasmic domain, promoting its degradation in the endoplasmic reticulum and preventing surface expression, aiding viral immune evasion. CD4 plays a central role in T-cell activation

and HIV infection and is a key focus in immunological and virological research.

PRODUCT DATA



Human CD4 Protein (C-His-Avi) was biotinylated in vitro using BirA ligase. SDS-PAGE analysis under non-reducing (P-) conditions shows the protein has a purity greater than 95%. A gel shift assay using co-incubation with streptavidin indicates that the biotinylation efficiency of the CD4 protein exceeds 90%.

This product is supplied subject to the terms and conditions at www.innocyto.com/web/terms.php and may only be used as provided in the stated terms. Products are for Research Use Only.