

## PE Human GITR (TNFRSF18) (C-His)

<b>Catalog Number:</b>	811501, 811502
<b>Size:</b>	25 ug, 100 ug
<b>Target Name:</b>	TNFRSF18, AITR, GITR, CD357
<b>Regulatory Status:</b>	RUO

### PRODUCT DETAILS

---

<b>Application:</b>	Flow Cytometry
<b>Format:</b>	Liquid, PE
<b>Expression Host:</b>	CHO
<b>Species:</b>	Human
<b>Sources:</b>	Recombinant Human Human GITR/TNFRSF18 (Gln26-Glu161) with C-terminus His-tag is expressed in CHO cell and conjugated to PE.
<b>Accession Number:</b>	Q9Y5U5
<b>Molecular Weight:</b>	The protein has a predicted molecular weight of 16.1 kDa. Under DTT-reducing conditions, it migrates at approximately 25 kDa on SDS-PAGE prior to conjugation.
<b>Affinity Tag:</b>	C-His
<b>Formulation:</b>	1xPBS buffer, pH7.4, 0.09% NaN3 with a carrier protein
<b>Endotoxin level:</b>	Not tested
<b>Protein Concentration:</b>	25µg size is bottled at 0.1mg/mL concentration. 100 µg size is bottled at lot specific concentration.
<b>Storage and Handling:</b>	Briefly centrifuge the vial upon receipt. An unopened vial may be stored at 2-8°C for up to six months.

### BACKGROUND INFORMATION

---

GITR (glucocorticoid-induced TNFR-related protein), also known as TNFRSF18 or CD357, is a 25 kD member of the TNF receptor superfamily that acts as the receptor for TNFSF18 (GITRL). It is primarily expressed on activated T cells and regulatory T cells and is upregulated upon T cell receptor engagement. GITR plays a key role in immune regulation by influencing T cell proliferation, TCR-mediated apoptosis, and the function of regulatory T cells, thereby contributing to the maintenance of self-tolerance. GITR signaling activates NF-κB via the TRAF2/NIK pathway and interacts with TRAF1-3. It is also implicated in T cell-endothelial cell interactions and the pathogenesis of autoimmune diseases.