

## APC Human GITR (TNFRSF18) (C-Fc)

|                           |                             |
|---------------------------|-----------------------------|
| <b>Catalog Number:</b>    | 811803, 811804              |
| <b>Size:</b>              | 25 ug, 100 ug               |
| <b>Target Name:</b>       | TNFRSF18, AITR, GITR, CD357 |
| <b>Regulatory Status:</b> | RUO                         |

### PRODUCT DETAILS

---

|                               |   |
|-------------------------------|---|
| <b>Application:</b>           | Flow Cytometry  |
| <b>Format:</b>                | Liquid, APC   |
| <b>Expression Host:</b>       | CHO   |
| <b>Species:</b>               | Human   |
| <b>Sources:</b>               | Recombinant Human GITR/TNFRSF18 (Gln26-Glu161) with C-terminus Fc-tag is expressed in CHO cell and conjugated to APC.   |
| <b>Accession Number:</b>      | Q9Y5U5  |
| <b>Molecular Weight:</b>      | The protein has a predicted molecular weight of 40.8 kDa. Under DTT-reducing conditions, it migrates at approximately 45-50 kDa on SDS-PAGE prior to conjugation. |
| <b>Affinity Tag:</b>          | C-Fc  |
| <b>Formulation:</b>           | 1xPBS buffer, pH7.4, 0.09% NaN3 with a carrier protein  |
| <b>Endotoxin level:</b>       | Not tested  |
| <b>Protein Concentration:</b> | 25µg size is bottled at 0.1mg/mL concentration. 100 µg size is bottled at lot specific concentration.   |
| <b>Storage and Handling:</b>  | Briefly centrifuge the vial upon receipt. An unopened vial may be stored at 2-8°C for up to six months.   |

### BACKGROUND INFORMATION

---

GITR (glucocorticoid-induced TNFR-related protein), also known as TNFRSF18 or CD357, is a 25 kD member of the TNF receptor superfamily that acts as the receptor for TNFSF18 (GITRL). It is primarily expressed on activated T cells and regulatory T cells and is upregulated upon T cell receptor engagement. GITR plays a key role in immune regulation by influencing T cell proliferation, TCR-mediated apoptosis, and the function of regulatory T cells, thereby contributing to the maintenance of self-tolerance. GITR signaling activates NF-κB via the TRAF2/NIK pathway and interacts with TRAF1-3. It is also implicated in T cell-endothelial cell interactions and the pathogenesis of autoimmune diseases.