

Human CD25 (IL-2R) Protein (C-Fc-Avi)

Catalog Number:	812801, 812802
Size:	25 ug, 100 ug
Target Name:	IL2RA, CD25, p55
Regulatory Status:	RUO


PRODUCT DETAILS

Application:	ELISA, BLI
Format:	Liquid, Purified
Expression Host:	CHO
Species:	Human
Sources:	Recombinant Human CD25 (Glu22-Cys213) with C-terminus Fc-Avi-tag is expressed in CHO cell.
Accession Number:	P01589
Molecular Weight:	The protein has a predicted molecular weight of 50.2 kDa. Under DTT-reducing conditions, it migrates at approximately 65 kDa on SDS-PAGE.
Affinity Tag:	C-Fc-Avi
Purity:	>95% based on SDS-PAGE under reducing condition
Formulation:	1xPBS buffer, pH7.4, 0.22 µm filtered
Endotoxin level:	Not tested
Protein Concentration:	25µg size is bottled at 0.2mg/mL concentration. 100 µg size is supplied at a lot-specific concentration.
Storage and Handling:	Briefly centrifuge the vial upon receipt. An unopened vial can be stored at 4°C for up to 2 weeks, or at -20°C or below for up to six months. The protein may be further diluted to 0.1 mg/mL using 0.22 µm-filtered PBS buffer (pH 7.4). For long-term storage, the diluted stock solution should be aliquoted and stored at ≤ -70°C to minimize freeze-thaw cycles. If additional dilution is required, carrier proteins such as FBS or BSA should be added to maintain protein stability.

BACKGROUND INFORMATION

CD25 (IL2RA), a type I transmembrane glycoprotein, is expressed on activated T cells, regulatory T cells, and dendritic cells. It binds IL-2 with low affinity on its own, but upon associating with the IL-2 receptor beta (CD122) and common gamma chain (CD132), it forms a high-affinity complex that triggers intracellular signaling pathways like MAPK and JAK/STAT. CD25 is involved in immune regulation and plays a role in autoimmune diseases, including type 1 diabetes. Soluble CD25 (sIL-2Ralpha) is released during immune responses and is a marker for tumor burden in hematological malignancies, including leukemias and lymphomas, as well as some solid tumors like melanoma and carcinomas. The presence of sIL-2Ralpha in circulation reflects tumor activity and is associated with immune modulation in cancer progression.

PRODUCT DATA



Human CD25 (C-Fc-Avi) on SDS-PAGE under reducing condition (P+) and non-reducing condition (P-). The gel was stained for 1 hour with BlinkBlue (catalog 700102). The purity of this protein appears to be greater than 95%.

This product is supplied subject to the terms and conditions at www.innocyto.com/web/terms.php and may only be used as provided in the stated terms. Products are for Research Use Only.