

PE Human CD25 (IL-2R α) Protein (C-Fc)

Catalog Number:	813001, 813002
Size:	25 ug, 100 ug
Target Name:	IL2RA, CD25, p55
Regulatory Status:	RUO

PRODUCT DETAILS

Application:	Flow Cytometry
Format:	Liquid, PE
Expression Host:	CHO
Species:	Human
Sources:	Recombinant Human CD25 (Glu22-Cys213) with C-terminus Fc-tag is expressed in CHO cell and conjugated to PE.
Accession Number:	P01589
Molecular Weight:	The protein has a predicted molecular weight of 48.0 kDa. Under DTT-reducing conditions, it migrates at approximately 65 kDa on SDS-PAGE prior to conjugation.
Affinity Tag:	C-Fc
Formulation:	1xPBS buffer, pH7.4, 0.09% NaN ₃ with a carrier protein
Endotoxin level:	Not tested
Protein Concentration:	25 μ g size is bottled at 0.1mg/mL concentration. 100 μ g size is bottled at lot specific concentration.
Storage and Handling:	Briefly centrifuge the vial upon receipt. An unopened vial may be stored at 2-8°C for up to six months.

BACKGROUND INFORMATION

CD25 (IL2RA), a type I transmembrane glycoprotein, is expressed on activated T cells, regulatory T cells, and dendritic cells. It binds IL-2 with low affinity on its own, but upon associating with the IL-2 receptor beta (CD122) and common gamma chain (CD132), it forms a high-affinity complex that triggers intracellular signaling pathways like MAPK and JAK/STAT. CD25 is involved in immune regulation and plays a role in autoimmune diseases, including type 1 diabetes. Soluble CD25 (sIL-2R α) is released during immune responses and is a marker for tumor burden in hematological malignancies, including leukemias and lymphomas, as well as some solid tumors like melanoma and carcinomas. The presence of sIL-2R α in circulation reflects tumor activity and is associated with immune modulation in cancer progression.