

## Human PD-L2 (CD273) Protein (C-Fc-Avi)

<b>Catalog Number:</b>	813401, 813402
<b>Size:</b>	25 ug, 100 ug
<b>Target Name:</b>	PDL2, , Butyrophilin B7-DC, CD273, PDCD1 ligand 2
<b>Regulatory Status:</b>	RUO

### PRODUCT DETAILS

---

<b>Application:</b>	ELISA, BLI
<b>Format:</b>	Liquid, Purified
<b>Expression Host:</b>	CHO
<b>Species:</b>	Human
<b>Sources:</b>	Recombinant Human PD-L2 (Leu20-Pro219) with C-terminus Fc-Avi-tag is expressed in CHO cell.
<b>Accession Number:</b>	Q9BQ51
<b>Molecular Weight:</b>	The protein has a predicted molecular weight of 51.2 kDa. Under DTT-reducing conditions, it migrates at approximately 65-75 kDa on SDS-PAGE.
<b>Affinity Tag:</b>	C-Fc-Avi
<b>Purity:</b>	>95% based on SDS-PAGE under reducing condition
<b>Formulation:</b>	1xPBS buffer, pH7.4, 0.22 µm filtered
<b>Endotoxin level:</b>	Not tested
<b>Protein Concentration:</b>	25µg size is bottled at 0.2mg/mL concentration. 100 µg size is supplied at a lot-specific concentration.
<b>Storage and Handling:</b>	Briefly centrifuge the vial upon receipt. An unopened vial can be stored at 4°C for up to 2 weeks, or at -20°C or below for up to six months. The protein may be further diluted to 0.1 mg/mL using 0.22 µm-filtered PBS buffer (pH 7.4). For long-term storage, the diluted stock solution should be aliquoted and stored at ≤ -70°C to minimize freeze-thaw cycles. If additional dilution is required, carrier proteins such as FBS or BSA should be added to maintain protein stability.

### BACKGROUND INFORMATION


---

Programmed death ligand 2 (PD-L2, also known as B7-DC or CD273) is a type I transmembrane protein and a member of the B7 family, containing one V-like and one C-like Ig domain in its extracellular region. It is mainly expressed on antigen-presenting cells, such as dendritic cells, macrophages, and B cells, and its expression can be induced by IFN-γ or LPS. PD-L2 serves as a high-affinity ligand for PD-1, a receptor expressed on activated T and B cells, and inhibits T cell activation, proliferation, and cytokine production, thus contributing to immune tolerance and evasion, particularly in tumors and autoimmunity. PD-L2 also plays a PD-1-independent role in asthma by regulating airway hyperresponsiveness through IL-12 production and binding to an alternative receptor, RGMb, involved in respiratory immune regulation. Its dual functions in both PD-1-dependent and independent pathways highlight PD-L2 as

a critical immunoregulatory molecule and a potential therapeutic target in cancer, autoimmunity, and allergic diseases.

## PRODUCT DATA

---



Human PD-L2 (C-Fc-Avi) on SDS-PAGE under reducing condition (P+) and non-reducing condition (P-). The gel was stained for 1 hour with BlinkBlue (catalog 700102). The purity of this protein appears to be greater than 95%.

This product is supplied subject to the terms and conditions at [www.innocyto.com/web/terms.php](http://www.innocyto.com/web/terms.php) and may only be used as provided in the stated terms. Products are for Research Use Only.