

## Biotin Human Vista Protein (C-Fc-Avi)

<b>Catalog Number:</b>	814003, 814004
<b>Size:</b>	25 ug, 100 ug
<b>Target Name:</b>	B7-H5, SISP1, Gi24, VISTA
<b>Regulatory Status:</b>	RUO

### PRODUCT DETAILS

---

<b>Application:</b>	ELISA, BLI
<b>Format:</b>	Liquid, Biotinylated
<b>Expression Host:</b>	CHO
<b>Species:</b>	Human
<b>Sources:</b>	Recombinant Human Vista (Phe33-Ala194) with C-terminus Fc-Avi-tag is expressed in CHO cell. This protein was site-specifically labeled with Biotin by BirA ligase.
<b>Accession Number:</b>	Q9H7M9
<b>Molecular Weight:</b>	The protein has a predicted molecular weight of 47 kDa. Under DTT-reducing conditions, it migrates at approximately 55-65 kDa on SDS-PAGE.
<b>Affinity Tag:</b>	C-Fc-Avi
<b>Purity:</b>	>95% based on SDS-PAGE under reducing condition
<b>Formulation:</b>	1xPBS buffer, pH7.4, 0.22 µm filtered
<b>Endotoxin level:</b>	Not tested
<b>Protein Concentration:</b>	25µg size is bottled at 0.2mg/mL concentration. 100 µg size is supplied at a lot-specific concentration.
<b>Storage and Handling:</b>	Briefly centrifuge the vial upon receipt. An unopened vial can be stored at 4°C for up to 2 weeks, or at -20°C or below for up to six months. The protein may be further diluted to 0.1 mg/mL using 0.22 µm-filtered PBS buffer (pH 7.4). For long-term storage, the diluted stock solution should be aliquoted and stored at ≤ -70°C to minimize freeze-thaw cycles. If additional dilution is required, carrier proteins such as FBS or BSA should be added to maintain protein stability.

### BACKGROUND INFORMATION


---

VISTA (V-domain Ig suppressor of T cell activation), also known as VSIR, PD-1H, or C10orf54, is a type I transmembrane protein with a single IgV-like extracellular domain and functions as an inhibitory immune checkpoint molecule. It is broadly expressed on myeloid cells, T cells, dendritic cells, and in tissues such as spleen and bone marrow. VISTA suppresses CD4<sup>+</sup> and CD8<sup>+</sup> T cell activation and cytokine production, contributing to immune tolerance and regulation. It may also play a role in embryonic stem cell differentiation by modulating BMP4 signaling. VISTA undergoes proteolytic cleavage, generating both soluble and membrane-bound fragments, and its interaction with PSGL1 in low-pH tumor microenvironments has been reported. Due to its immunosuppressive

properties, VISTA is considered a promising target for cancer immunotherapy and may be involved in inflammatory conditions such as chronic rhinosinusitis with nasal polyps.

## PRODUCT DATA

---



Human Vista (C-Fc-Avi) was biotinylated in vitro using BirA ligase. SDS-PAGE analysis under reducing (P+) and non-reducing (P-) conditions shows the protein has a purity greater than 95%. A gel shift assay using co-incubation with streptavidin indicates that the biotinylation efficiency of the Vista protein exceeds 80%.

This product is supplied subject to the terms and conditions at [www.innocyto.com/web/terms.php](http://www.innocyto.com/web/terms.php) and may only be used as provided in the stated terms. Products are for Research Use Only.