www.innocyto.com

InnoCyto Inc.

15375 Barranca Pkwy, Suite I-103
Irvine, CA 92618
Email: info@innocyto.com
Phone: 1-949-418-7543

APC Human CD200 (OX-2) Protein (C-His)

Catalog Number:	814803, 814804
Size:	25 ug, 100 ug
Target Name:	CD200, MOX1, MOX2, MRC, OX-2, My033
Regulatory Status:	RUO

PRODUCT DETAILS

Application:	Flow Cytometry
Format:	Liquid, APC
Expression Host:	CHO
Species:	Human
Sources:	Recombinant Human CD200 (Gln31-Gly232) with C-terminus His-tag is expressed in CHO cell and conjugated to APC.
Accession Number:	P41217
Molecular Weight:	The protein has a predicted molecular weight of 24.1 kDa. Under DTT-reducing conditions, it migrates at approximately 35-45 kDa on SDS-PAGE prior to conjugation.
Affinity Tag:	C-His
Formulation:	1xPBS buffer, pH7.4, 0.09% NaN3 with a carrier protein
Endotoxin level:	Not tested
Protein Concentration:	25μg size is bottled at 0.1mg/mL concentration. 100 μg size is bottled at lot specific concentration.
Storage and Handling:	Briefly centrifuge the vial upon receipt. An unopened vial may be stored at 2–8°C for up to six months.

BACKGROUND INFORMATION

CD200 (OX-2) is a cell surface glycoprotein that regulates immune responses through its receptor, CD200R, mainly expressed on myeloid cells. It plays a crucial role in suppressing alloimmune and autoimmune responses, contributing to immune privilege in various tissues. CD200-CD200R signaling is essential in regulating anti-tumor immunity, with overexpression linked to malignancies like chronic lymphocytic leukemia (CLL) and cancer stem cells. Additionally, CD200 signaling is involved in the central nervous system, particularly in diseases like Parkinson's, where it affects microglia activation. Elevated CD200 expression is associated with reduced transplant rejection, autoimmunity, and allergic diseases, but may also promote tumor cell survival. While it helps prevent graft rejection and autoimmune diseases, high CD200 levels in cancers have been linked to poor prognosis. This dual role underscores the complexity of CD200 in immune regulation and its potential as both a therapeutic target and a biomarker.