

Human GM2A Protein (C-FLAG)

| | |
|---------------------------|---------------------------------|
| Catalog Number: | 601601, 601602 |
| Size: | 25 ug, 100 ug |
| Target Name: | GM2A, Ganglioside GM2 activator |
| Regulatory Status: | RUO |


PRODUCT DETAILS

| | |
|-------------------------------|--|
| Application: | ELISA, BLI |
| Format: | Liquid, Purified |
| Expression Host: | HEK293 |
| Species: | Human |
| Accession Number: | P17900 |
| Sources: | Recombinant human GM2A (Ser32-Ile193) with C-terminus DYKDDDDK tag was expressed in 293 cells. |
| Molecular Weight: | This protein has a predicted molecular weight of 18.9 kDa. Under DTT-reducing conditions, the protein migrates at approximately 20 kDa on SDS-PAGE. |
| Affinity Tag: | C-DYKDDDDK |
| Purity: | >95% based on SDS-PAGE under reducing condition |
| Formulation: | 1xPBS with 300mM NaCl, pH7.4, 0.22 μ m filtered |
| Endotoxin level: | Not tested |
| Protein Concentration: | 25 μ g size is bottled at 0.2mg/mL concentration. 100 μ g size is supplied at a lot-specific concentration. |
| Storage and Handling: | Briefly centrifuge the vial upon receipt. An unopened vial can be stored at 4°C for up to 2 weeks, or at -20°C or below for up to six months. The protein may be further diluted to 0.1 mg/mL using 0.22 μ m-filtered PBS buffer (pH 7.4). For long-term storage, the diluted stock solution should be aliquoted and stored at \leq -70°C to minimize freeze-thaw cycles. If additional dilution is required, carrier proteins such as FBS or BSA should be added to maintain protein stability. |

BACKGROUND INFORMATION

GM2 ganglioside activator (GM2A) is a lipid transfer protein of the ML domain family that facilitates the degradation of ganglioside GM2 by acting as a substrate-specific cofactor for β -hexosaminidase A. GM2A extracts GM2 from membranes and presents it in soluble form to β -hexosaminidase A for cleavage, enabling conversion to GM3. It can bind several single-chain phospholipids and fatty acids and shows some calcium-independent phospholipase activity. Mutations in GM2A cause GM2-gangliosidosis type AB, a Tay-Sachs disease variant characterized by impaired GM2 degradation and lipid accumulation.

PRODUCT DATA



Purified Human GM2A (His24-Ile193, with C-terminus DYKDDDDK tag, 293 expressed) final product on SDS-PAGE under reducing (P+) conditions. The purity of Human GM2A appears to be greater than 95%.