

## SARS-CoV-2 Spike RBD (N501Y) Protein (C-His-Avi)

<b>Catalog Number:</b>	602801, 602802
<b>Size:</b>	25 ug, 100 ug
<b>Target Name:</b>	SARS-CoV2 RBD, Spike RBD Protein, RBD Protein
<b>Regulatory Status:</b>	RUO

### PRODUCT DETAILS

---

<b>Application:</b>	ELISA, BLI
<b>Format:</b>	Liquid, Purified
<b>Expression Host:</b>	HEK293
<b>Species:</b>	SARS-CoV-2
<b>Accession Number:</b>	QHD43416.1
<b>Sources:</b>	Recombinant SARS-CoV-2 S protein RBD (Arg319-Phe541, N501Y) with C-terminus His-Avi tag was expressed in 293 Cells.
<b>Molecular Weight:</b>	This protein has a predicted molecular weight of 29 kDa. Under DTT-reducing conditions, the protein migrates at approximately 30-40 kDa on SDS-PAGE.
<b>Affinity Tag:</b>	C-His-Avi
<b>Purity:</b>	>95% based on SDS-PAGE under reducing condition
<b>Formulation:</b>	1xPBS buffer, pH7.4, 0.22 µm filtered
<b>Endotoxin level:</b>	Not tested
<b>Protein Concentration:</b>	25µg size is bottled at 0.2mg/mL concentration. 100 µg size is supplied at a lot-specific concentration.
<b>Storage and Handling:</b>	Briefly centrifuge the vial upon receipt. An unopened vial can be stored at 4°C for up to 2 weeks, or at -20°C or below for up to six months. The protein may be further diluted to 0.1 mg/mL using 0.22 µm-filtered PBS buffer (pH 7.4). For long-term storage, the diluted stock solution should be aliquoted and stored at ≤ -70°C to minimize freeze-thaw cycles. If additional dilution is required, carrier proteins such as FBS or BSA should be added to maintain protein stability.


### BACKGROUND INFORMATION

---

The receptor-binding domain (RBD) of SARS-CoV-2 is a critical region of the spike (S) protein that directly interacts with the human ACE2 receptor to facilitate viral entry into host cells. It is located within the S1 subunit of the spike protein and adopts a compact, globular structure stabilized by disulfide bonds. Due to its essential role in viral entry, the RBD is a primary target for neutralizing antibodies, vaccines, and therapeutic interventions. Mutations in the RBD—seen in various SARS-CoV-2 variants—can increase binding affinity to ACE2 or help evade immune responses. The N501Y mutation enhances the virus's ability to bind to the ACE2 receptor, likely contributing to increased transmissibility.

**PRODUCT DATA**

---



SARS-CoV2 RBD (N501Y) Protein with C-His-Avi tag on SDS-PAGE under reducing conditions. The gel was stained for 1 hour with BlinkBlue Protein Staining Buffer (Catalog 700102). The purity of this protein appears to be greater than 95%.

This product is supplied subject to the terms and conditions at [www.innocyto.com/web/terms.php](http://www.innocyto.com/web/terms.php) and may only be used as provided in the stated terms. Products are for Research Use Only.