

## Biotin Human GITRL/TNFSF18 (N-His-Avi)

<b>Catalog Number:</b>	606603, 606604
<b>Size:</b>	25 ug, 100 ug
<b>Target Name:</b>	TNFSF18, GITR Ligand, AITRL, TL6, GITRL
<b>Regulatory Status:</b>	RUO

### PRODUCT DETAILS

---

<b>Application:</b>	ELISA, BLI
<b>Format:</b>	Liquid, Biotinylated
<b>Expression Host:</b>	CHO
<b>Species:</b>	Human
<b>Accession Number:</b>	Q9UNG2
<b>Sources:</b>	Recombinant Human TNFSF18 (Glu52-Ser177) with N-terminus His-Avi tag is expressed in CHO cells. This protein was site-specifically labeled with Biotin by BirA ligase.
<b>Molecular Weight:</b>	This protein has the predicted molecular weight of 18.3 kD. Under DTT-reducing conditions, the protein migrates at approximately 20 kD on SDS-PAGE
<b>Affinity Tag:</b>	N-His-Avi
<b>Purity:</b>	>95% based on SDS-PAGE under reducing condition
<b>Formulation:</b>	1xPBS buffer, pH7.4, 0.22 µm filtered
<b>Endotoxin level:</b>	Less than 0.1 EU/µg protein as determined by the LAL method
<b>Protein Concentration:</b>	25µg size is bottled at 0.2mg/mL concentration. 100 µg size is supplied at a lot-specific concentration.
<b>Storage and Handling:</b>	Briefly centrifuge the vial upon receipt. An unopened vial can be stored at 4°C for up to 2 weeks, or at -20°C or below for up to six months. The protein may be further diluted to 0.1 mg/mL using 0.22 µm-filtered PBS buffer (pH 7.4). For long-term storage, the diluted stock solution should be aliquoted and stored at ≤ -70°C to minimize freeze-thaw cycles. If additional dilution is required, carrier proteins such as FBS or BSA should be added to maintain protein stability.


### BACKGROUND INFORMATION

---

Human GITRL (Glucocorticoid-Induced TNFR-Related Ligand), also known as TNFSF18, is a member of the tumor necrosis factor (TNF) superfamily. It is a type II transmembrane protein primarily expressed on activated antigen-presenting cells, including B cells, dendritic cells, and macrophages. GITRL binds to its receptor GITR, which is found on activated T cells, regulatory T cells (Tregs), and natural killer (NK) cells. The GITR-GITRL interaction delivers a co-stimulatory signal that promotes T cell activation, proliferation, and survival, while also modulating the suppressive activity of Tregs. This signaling pathway plays a key role in immune regulation, inflammation, and antitumor immunity.

## PRODUCT DATA

---



Human GITRL (TNFSF18) Protein (N-His-Avi) was biotinylated in vitro using BirA ligase. SDS-PAGE analysis under reducing (P-) conditions shows the protein has a purity greater than 95%. A gel shift assay using co-incubation with streptavidin indicates that the biotinylation efficiency of the GITRL protein exceeds 70%.

This product is supplied subject to the terms and conditions at [www.innocyto.com/web/terms.php](http://www.innocyto.com/web/terms.php) and may only be used as provided in the stated terms. Products are for Research Use Only.