

Anti-Mouse Ly-6A/E (Sca-1) Antibody

Catalog Number:	200601, 200602
Size:	100 ug, 500 ug
Target Name:	Ly-6A/E, Sca-1
Regulatory Status:	RUO

PRODUCT DETAILS

Clone:	D7
Application:	Flow Cytometry
Reactivity:	Mouse
Format:	Purified
Isotype:	Rat IgG2a
Antibody Type:	Monoclonal
Formulation:	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide
Protein Concentration:	0.5 mg/mL
Storage&Handling:	The antibody solution should be stored between 2°C and 8°C
Recommended Usage:	For flow cytometric staining, it is recommended to use less than 0.2 µg of this reagent per 0.5-1.0 million cells in a 100 µL volume. Optimal reagent performance should be determined by titration for each specific application.
Isotype Control:	300201

BACKGROUND INFORMATION

Ly-6A/E, commonly known as stem cell antigen-1 (Sca-1), is a glycosylphosphatidylinositol (GPI)-anchored cell surface protein that serves as an important marker for murine hematopoietic stem cells and other progenitor populations. It belongs to the Ly-6/uPAR superfamily, which consists of small, cysteine-rich proteins involved in cell signaling, adhesion, and immune regulation. Functionally, Sca-1 contributes to stem cell maintenance, tissue regeneration, and immune responses, making it a key molecule for studying stem cell biology and immunoregulation in mice.

Structurally, Ly-6A/E is a relatively small protein of about 18-20 kDa, characterized by a conserved “three-finger” Ly-6 domain stabilized by multiple intramolecular disulfide bonds. It is tethered to the cell membrane via a GPI anchor rather than a transmembrane domain, allowing lateral mobility within the lipid bilayer and association with signaling complexes. Sca-1 lacks intrinsic signaling motifs but modulates cell function through its interactions with other membrane-associated proteins, influencing downstream signaling cascades involving Src-family kinases and MAPK pathways.


While the precise ligands of Sca-1 remain not fully defined, its activity is thought to depend on interactions with lipid raft-associated coreceptors and possibly extracellular matrix components. Sca-1 expression is dynamically regulated by cytokines and growth factors, including interferons, which enhance its expression in both stem and immune cells. Sca-1 plays multiple physiological

roles. In the hematopoietic system, it marks multipotent progenitors within the bone marrow and contributes to self-renewal and differentiation capacity. In non-hematopoietic tissues, such as skeletal muscle and the heart, Sca-1 marks resident stem or progenitor cells involved in tissue repair following injury. Within the immune system, it is expressed on activated T cells and influences T cell activation and cytokine production.

In disease, altered Sca-1 expression or function has been associated with impaired tissue regeneration, inflammatory conditions, and tumor progression. Sca-1-positive cells have been identified in various cancers, where they may contribute to tumor initiation, metastasis, and resistance to therapy, acting as markers of tumor-propagating cell populations.

Therapeutically, Ly-6A/E serves as a valuable marker for isolating stem and progenitor cells for transplantation and regenerative medicine. Understanding its regulatory role in stem cell signaling could improve strategies for tissue repair and immunotherapy. Additionally, Sca-1-associated pathways may offer targets for modulating cancer stem cell behavior and enhancing therapeutic response in experimental disease models.

PRODUCT DATA



Mouse splenocytes stained with either Anti-Mouse Ly-6A/E (Sca-1) clone D7 (blue histogram) or an isotype control (gray histogram), followed by PE conjugated anti-rat IgG.

This product is supplied subject to the terms and conditions at www.innocyto.com/web/terms.php and may only be used as provided in the stated terms. Products are for Research Use Only.