

APC Anti-Human TCR Vb13.1 Antibody

Catalog Number:	106311, 106312
Size:	25 tests, 100 tests
Target Name:	TCR Vb13.1, T cell receptor V β 13.1 chain, TCRBV13.1, TCRBV13
Regulatory Status:	RUO


PRODUCT DETAILS

Clone:	H131
Application:	Flow Cytometry
Reactivity:	Human
Format:	APC
Isotype:	Mouse IgG2b
Antibody Type:	Monoclonal
Formulation:	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 0.2% (w/v) BSA
Protein Concentration:	Supplied at a lot-specific concentration.
Storage&Handling:	The antibody solution should be stored undiluted between 2°C and 8°C, and protected from prolonged exposure to light. Do not freeze.
Recommended Usage:	For flow cytometric staining, it is recommended to use 5 μ L of this reagent per 0.5-1.0 million cells in a 100 μ L volume. Optimal reagent performance should be determined by titration for each specific application. APC has an excitation max at 650 nm and an emission max at 660 nm.
Excitation Laser:	Red Laser (633 nm)
Isotype Control:	301621
RRID:	AB_3738718

BACKGROUND INFORMATION

TCR V β 13.1 is a variant of the TCR β chain. The receptor is complexed with the TCR α chain and belongs to the immunoglobulin superfamily. It is expressed on a subset of T cells and some T cell clones. Variability in the β chain is generated by V β , D β , and J β gene rearrangement, while variability in the α chain is generated by V α and J α rearrangement. TCR V β 13.1 has been shown to be related to antigen recognition and inflammation as well as diseases, such as HIV and multiple sclerosis.

PRODUCT DATA



Human peripheral blood lymphocytes stained iF488 anti-human CD3 and APC Anti-Human TCR Vb13.1 clone H131 (right panel) or APC isotype control (left panel).

This product is supplied subject to the terms and conditions at www.innocyto.com/web/terms.php and may only be used as provided in the stated terms. Products are for Research Use Only.