

## APC Anti-Human CD298 Antibody

|                           |  |
|---------------------------|--|
| <b>Catalog Number:</b>    | 109111, 109112                                 |
| <b>Size:</b>              | 25 tests, 100 tests                            |
| <b>Target Name:</b>       | CD298, ATPB-3, Na, K-ATPase beta-3 polypeptide |
| <b>Regulatory Status:</b> | RUO  |

### PRODUCT DETAILS

---

|                               |   |
|-------------------------------|---|
| <b>Clone:</b>                 | LNH-94  |
| <b>Application:</b>           | Flow Cytometry  |
| <b>Reactivity:</b>            | Human   |
| <b>Format:</b>                | APC   |
| <b>Isotype:</b>               | Mouse IgG1  |
| <b>Antibody Type:</b>         | Monoclonal  |
| <b>Formulation:</b>           | Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 0.2% (w/v) BSA   |
| <b>Protein Concentration:</b> | Supplied at a lot-specific concentration.   |
| <b>Storage&amp;Handling:</b>  | The antibody solution should be stored undiluted between 2°C and 8°C, and protected from prolonged exposure to light. Do not freeze.  |
| <b>Recommended Usage:</b>     | For flow cytometric staining, it is recommended to use 5 µL of this reagent per 0.5-1.0 million cells in a 100 µL volume. Optimal reagent performance should be determined by titration for each specific application. APC has an excitation max at 650 nm and an emission max at 660 nm. |
| <b>Excitation Laser:</b>      | Red Laser (633 nm)  |
| <b>Isotype Control:</b>       | 301403  |

### BACKGROUND INFORMATION

---

CD298, also known as the sodium/potassium-transporting ATPase subunit beta-3 or ATP1B3, is a 42 kDa type II transmembrane glycoprotein encoded by the ATP1B3 gene in humans. It is part of the Na<sup>+</sup>/K<sup>+</sup>-ATPase enzyme complex, which consists of a catalytic  $\alpha$  subunit and a regulatory  $\beta$  subunit. CD298 belongs to the family of Na<sup>+</sup>/K<sup>+</sup> and H<sup>+</sup>/K<sup>+</sup> ATPase beta chain proteins and specifically to the Na<sup>+</sup>/K<sup>+</sup>-ATPase subfamily. This integral plasma membrane protein is essential for establishing and maintaining the electrochemical gradients of sodium and potassium ions across the cell membrane—gradients that are crucial for osmoregulation, sodium-coupled transport of various molecules, and the electrical excitability of nerve and muscle cells. The  $\beta$ 3 subunit plays a regulatory role by assisting in the proper assembly and membrane localization of the  $\alpha/\beta$  heterodimer complex. CD298 is broadly expressed across various tissues, including all leukocyte populations.

### PRODUCT DATA

---



Human peripheral blood lymphocytes stained either APC Anti-Human CD298 clone LNH-94 (color-filled histogram) or an isotype control (gray histogram).

This product is supplied subject to the terms and conditions at [www.innocyto.com/web/terms.php](http://www.innocyto.com/web/terms.php) and may only be used as provided in the stated terms. Products are for Research Use Only.