

## Technical Data Sheet

### APC Anti-Mouse CD19

**Catalog Number:** 201513, 201514  
**Size:** 25 tests, 100 tests  
**Target Name:** CD19, B4, CVID3, Leu-12, MGC12802  
**Regulatory Status:** RUO

#### Product Details

---

**Clone:** 6D5  
**Application:** FC  
**Reactivity:** Mouse  
**Format:** APC  
**Isotype:** Rat IgG2a  
**Antibody Type:** Monoclonal  
**Formulation:** Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 0.2% (w/v) BSA  
**Protein Concentration:** Supplied at a lot-specific concentration.  
**Storage&Handling:** The antibody solution should be stored undiluted between 2°C and 8°C, and protected from prolonged exposure to light. Do not freeze.  
**Recommended Usage:** For flow cytometric staining, it is recommended to use 5 uL of this reagent per 0.5-1.0 million cells in a 100 µL volume. Optimal reagent performance should be determined by titration for each specific application.  
**Excitation Laser:** Red Laser (633 nm)  
**Release Date:** Nov-25


#### Background Information

---

CD19, also known as B-lymphocyte antigen CD19, B4, Leu-12, or CVID3, is a type I transmembrane glycoprotein encoded by the CD19 gene. It is expressed throughout B cell development from the pro-B stage to mature B cells and is present on most B lineage cells. CD19 functions as an adaptor protein, recruiting cytoplasmic signaling molecules to the membrane, and forms part of the CD19/CD21/CD81 complex, which lowers the activation threshold for B cell receptor (BCR) signaling. Because of its consistent expression on B cells, CD19 serves as a key biomarker for B cell development and is widely used in the diagnosis of B cell lymphomas and leukemias. It is also a major therapeutic target in immunotherapies, including CAR T-cell therapy for B cell malignancies.

#### Product Data

---



Mouse splenocytes were stained either APC Anti-Mouse CD19 clone 6D5 (color-filled histogram) or an isotype control (gray histogram).