

### InnoCyto Inc.

15375 Barranca Pkwy, Suite I-103 Irvine, CA 92618

## **Technical Data Sheet**

APC Anti-Human CD197

Catalog Number: 107313, 107314

Size: 25 tests, 100 tests

Target Name: CD197, CCR7, BLR2, CDw197, EBI1, CMKBR7

Regulatory Status: RUO

#### **Product Details**

Clone: 197AM2a Application: FC Reactivity: Human Format: APC

Isotype: Mouse IgG2a

Antibody Type: Monoclonal

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 0.2% (w/v)

BSA

**Protein Concentration:** Supplied at a lot-specific concentration.

Storage&Handling: The antibody solution should be stored undiluted between 2°C and 8°C, and

protected from prolonged exposure to light. Do not freeze.

**Recommended Usage:** For flow cytometric staining, it is recommended to use 5 uL of this reagent per 0.5-1.0 million cells in a 100 μL volume. Optimal reagent performance should be determined by

titration for each specific application. **Excitation Laser:** Red Laser (633 nm)

Release Date: Dec-25

#### **Background Information**


C-C chemokine receptor type 7 is a protein that in humans is encoded by the CCR7 gene. CD197 is a chemokine receptor that binds CCL19 and CCL21. CCR7 is expressed on naive T cells, regulatory T cell, memory T cell, dendritic cell, mature B cell. Through regulating the homing of CD197 expressing cells into secodory lymphoid organs, it affects the interaction among these cells and thus the immune respose.

#### **Product Data**




# InnoCyto Inc.

15375 Barranca Pkwy, Suite I-103 Irvine, CA 92618



Human peripheral blood lymphocytes was stained with APC Anti-Human CD197 clone 197AM2a (color-filled) or an isotype control (gray histogram).



Human peripheral blood lymphocytes was co-stained with APC Anti-Human CD197 clone 197AM2a and FITC Anti-Human CD3 (clone OKT3)