

## APC Anti-Human TCR Vb5.2/5.3 Antibody

|                           |                     |
|---------------------------|---------------------|
| <b>Catalog Number:</b>    | 107603, 107604      |
| <b>Size:</b>              | 25 tests, 100 tests |
| <b>Target Name:</b>       | TCR Vb5.2/5.3       |
| <b>Regulatory Status:</b> | RUO                 |

### PRODUCT DETAILS

---

|                               |   |
|-------------------------------|---|
| <b>Clone:</b>                 | MH3-2   |
| <b>Application:</b>           | Flow Cytometry  |
| <b>Reactivity:</b>            | Human   |
| <b>Format:</b>                | APC   |
| <b>Isotype:</b>               | Mouse IgG2a   |
| <b>Antibody Type:</b>         | Monoclonal  |
| <b>Formulation:</b>           | Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 0.2% (w/v) BSA   |
| <b>Protein Concentration:</b> | Supplied at a lot-specific concentration.   |
| <b>Storage&amp;Handling:</b>  | The antibody solution should be stored undiluted between 2°C and 8°C, and protected from prolonged exposure to light. Do not freeze.  |
| <b>Recommended Usage:</b>     | For flow cytometric staining, it is recommended to use 5 µL of this reagent per 0.5-1.0 million cells in a 100 µL volume. Optimal reagent performance should be determined by titration for each specific application. APC has an excitation max at 650 nm and an emission max at 660 nm. |
| <b>Excitation Laser:</b>      | Red Laser (633 nm)  |
| <b>Isotype Control:</b>       | 301503  |



### BACKGROUND INFORMATION

---

Vβ5 T cell receptor (TCR Vβ5) is a variant of the TCR β chain which is heterodimerized with TCR α chain and expressed on a subset of αβ+ T cells. TCR α/β recognizes peptides presented by the MHC molecules and initiates T cell mediated immune responses. Patients with multiple sclerosis present an expansion of Vβ5.2+/5.3+ autoreactive T cells.

### PRODUCT DATA

---



Human peripheral blood lymphocytes stained with FITC Anti-human CD3 and APC Anti-Human TCR Vb5.2/5.3 (TRbV28) clone MH3-2 (right panel). Human peripheral blood

lymphocytes stained was stained with APC Anti-Human TCR Vb5.2/5.3 (TRbV28) clone MH3-2 (color-filled histogram, left panel) .

This product is supplied subject to the terms and conditions at [www.innocyto.com/web/terms.php](http://www.innocyto.com/web/terms.php) and may only be used as provided in the stated terms. Products are for Research Use Only.