

InnoCyto Inc.

15375 Barranca Pkwy, Suite I-103 Irvine, CA 92618

Technical Data Sheet

PE/Cyanine7 Anti-Mouse/human CD45R/B220

Catalog Number: 200913, 200914

Size: 25 tests, 100 tests
Target Name: CD45R, B220
Regulatory Status: RUO

Product Details

Clone: RA3-6B2 Application: FC

Reactivity: Human, Mouse Format: PE/Cyanine7 Isotype: Rat IgG2a

Antibody Type: Monoclonal

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 0.2% (w/v)

BSA

Protein Concentration: Supplied at a lot-specific concentration.

Storage&Handling: The antibody solution should be stored undiluted between 2°C and 8°C, and

protected from prolonged exposure to light. Do not freeze.

Recommended Usage: For flow cytometric staining, it is recommended to use 5 uL of this reagent per 0.5-1.0 million cells in a 100 µL volume. Optimal reagent performance should be determined by titration for each specific application.

Excitation Laser: Blue Laser (488 nm) Green/Yellow laser (532/561nm)

Release Date: Dec-25

Background Information


CD45R, also known as B220, is an isoform of CD45 (PTPRC), a member of the protein tyrosine phosphatase (PTP) family. With a molecular weight of approximately 180–240 kDa, CD45R is a transmembrane glycoprotein expressed on B cells at all developmental stages, as well as on activated B cells, subsets of T and NK cells, and certain abnormal T cell populations involved in systemic autoimmunity in MRL-Faslpr and MRL-Fasgld mice. Structurally, CD45R consists of an extracellular domain, a single transmembrane region, and two tandem intracytoplasmic catalytic domains. Functionally, CD45R plays a critical role in T cell receptor (TCR) and B cell receptor (BCR) signaling by regulating Src-family kinases, which are essential for lymphocyte activation and development. The primary ligands for CD45 include galectin-1, CD2, CD3, and CD4. CD45R is widely used as a pan-B cell marker, though CD19 may provide greater specificity for B cells.

Product Data



InnoCyto Inc.

15375 Barranca Pkwy, Suite I-103 Irvine, CA 92618



Mouse splenocytes were stained with PE/Cyanine7 Anti-Human/Mouse CD45R/B220 clone RA3-6B2 (color-filled histogram).