

## Purified anti-mouse/human GL7 Antigen Antibody

<b>Catalog Number:</b>	112101, 112102
<b>Size:</b>	25 ug, 100 ug
<b>Target Name:</b>	T and B cell activation marker T and B cell activation marker, Ly77
<b>Regulatory Status:</b>	RUO

### PRODUCT DETAILS

---

<b>Clone:</b>	GL7-mM
<b>Application:</b>	Flow Cytometry
<b>Reactivity:</b>	Human, mouse
<b>Format:</b>	Purified
<b>Isotype:</b>	mouse IgM
<b>Antibody Type:</b>	Monoclonal
<b>Formulation:</b>	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide
<b>Protein Concentration:</b>	0.5 mg/mL
<b>Storage&amp;Handling:</b>	The antibody solution should be stored between 2°C and 8°C
<b>Isotype Control:</b>	301601

### BACKGROUND INFORMATION

---

**GL7 antigen** is a cell-surface carbohydrate epitope commonly used as a marker of activated B cells, particularly **germinal center (GC) B cells**. It was originally identified through the monoclonal antibody GL7, which recognizes a specific glycan structure expressed on lymphocytes during immune activation. In both mouse and human immunology research, GL7 expression is widely used to identify germinal center B cells in secondary lymphoid organs such as lymph nodes, spleen, and Peyer's patches. These cells are actively undergoing proliferation, somatic hypermutation, and affinity maturation during adaptive immune responses.

Structurally, GL7 is not a classical protein antigen but rather a **glycan epitope** present on glycoproteins and glycolipids at the cell surface. The epitope corresponds to a modified **sialylated lactosamine carbohydrate structure** that becomes exposed when certain sialic acid residues are enzymatically altered during B-cell activation. Because the GL7 determinant is carbohydrate-based, it may appear on multiple membrane proteins or lipids that carry the appropriate glycosylation pattern. This explains why GL7 is considered a differentiation marker rather than a receptor with a single defined protein backbone.

Unlike classical immune receptors, GL7 does not have a well-defined ligand or signaling function. Instead, it reflects changes in cellular glycosylation associated with activation and germinal center differentiation. The appearance of GL7 correlates with B-cell interactions with T follicular helper cells and antigen stimulation within germinal centers. As a result, GL7 is often used together with markers such as B220, CD95 (Fas), and IgD to identify germinal center B-cell populations by flow cytometry or immunohistochemistry.

GL7 expression is associated with immune responses and can also appear in pathological conditions involving abnormal germinal

center activity. For example, elevated germinal center reactions marked by GL7-positive B cells may occur in autoimmune diseases, chronic infections, or certain B-cell lymphomas. Although GL7 itself is not typically targeted therapeutically, it is widely used in research and preclinical studies to monitor germinal center responses, vaccine-induced B-cell activation, and immune dysregulation. In this way, GL7 serves as an important biomarker for studying adaptive immune responses and evaluating immunotherapies or vaccine efficacy.

This product is supplied subject to the terms and conditions at [www.innocyto.com/web/terms.php](http://www.innocyto.com/web/terms.php) and may only be used as provided in the stated terms. Products are for Research Use Only.