

Recombinant Human IL-12 Protein

Catalog Number:	630101, 630102
Size:	10 µg, 100 µg
Target Name:	IL-12, Natural killer cell stimulatory factor (NKSF), cytotoxic lymphocyte maturation factor (CLMF)
Regulatory Status:	RUO

PRODUCT DETAILS

Application:	Bioassay
Format:	Lyophilized from sterile PBS, pH 7.4.
Expression Host:	HEK293
Species:	Human
accession number:	P29459-1 & NP_002178.2
Sources:	A DNA sequence encoding the human IL12A (P29459-1)(Met1-Ser219) was expressed, constructed the plasmid 1; A DNA sequence encoding the human IL12B (NP_002178.2)(Met1-Ser328) was expressed, constructed the plasmid 2. The two plasmids were co-expressed and the human IL12A&IL12B heterodimer was purified.
Molecular Weight:	The recombinant heterodimer of human IL12A&IL12B comprises 503(197+306) amino acids and has a calculated molecular mass of 57.2(22.5+34.7) KDa. The apparent molecular mass of human IL12 heterodimer is approximately 34 & 41 kDa respectively in SDS-PAGE under reducing conditions.
Affinity Tag:	None
Purity:	≥ 95 % as determined by SDS-PAGE. ≥ 95 % as determined by SEC-HPLC.
Endotoxin level:	
Protein Concentration:	Lyophilized
Storage and Handling:	Proteins are stable for up to twelve months from date of receipt at -20°C to -80°C. Store it under sterile conditions at -20°C to -80°C. It is recommended that the protein be aliquoted for optimal storage. Avoid repeated freeze-thaw cycles.

BACKGROUND INFORMATION

Human interleukin-12 (IL-12) is a proinflammatory cytokine that plays a central role in bridging innate and adaptive immunity. It is primarily produced by antigen-presenting cells such as dendritic cells, macrophages, and B cells in response to pathogens. IL-12 promotes the differentiation of naïve CD4+ T cells into T helper 1 (Th1) cells and enhances the cytotoxic activity of natural killer (NK) cells and CD8+ T lymphocytes. A key function of IL-12 is the induction of interferon-gamma (IFN-γ) production, which amplifies cellular immune responses against intracellular pathogens.

Structurally, IL-12 is a heterodimer composed of two covalently linked subunits: p35 and p40. These subunits are encoded by

separate genes and must assemble to form the biologically active cytokine, often referred to as IL-12p70. IL-12 signals through a receptor complex consisting of IL-12R β 1 and IL-12R β 2 chains, leading to activation of the JAK-STAT signaling pathway, particularly STAT4.

In disease, dysregulated IL-12 production is associated with chronic inflammation and autoimmune conditions such as psoriasis and inflammatory bowel disease. Conversely, insufficient IL-12 activity can impair host defense against infections. Therapeutically, IL-12 has been explored as an immunostimulatory agent in cancer, aiming to boost anti-tumor immunity. However, systemic toxicity has limited its direct use, leading to alternative strategies such as targeted delivery or modulation of its signaling pathway.

This product is supplied subject to the terms and conditions at www.innocyto.com/web/terms.php and may only be used as provided in the stated terms. Products are for Research Use Only.