

iF700 Anti-Mouse CD16/32

Catalog Number:	201817, 201818
Size:	25 µg, 100 µg
Target Name:	CD16/32
Regulatory Status:	RUO


PRODUCT DETAILS

Clone:	2.4G2
Application:	Flow Cytometry
Reactivity:	Mouse
Format:	iF700
Isotype:	Rat IgG2b
Antibody Type:	Monoclonal
Formulation:	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 0.2% (w/v) BSA
Protein Concentration:	0.2 mg/mL
Storage & Handling:	The antibody solution should be stored undiluted between 2°C and 8°C, and protected from prolonged exposure to light. Do not freeze.
Recommended Usage:	For flow cytometric staining, it is recommended to use less than 0.2 µg of this reagent per 0.5-1.0 million cells in a 100 µL volume. Optimal reagent performance should be determined by titration for each specific application. iF700 has an excitation max at 690 nm and an emission max at 710 nm.
Excitation Laser:	Red Laser (633 nm)
Isotype Control:	303611

BACKGROUND INFORMATION

CD16/32 are Fc-gamma receptors (FcγRs) expressed on a variety of immune cells, including B cells, monocytes/macrophages, NK cells, granulocytes, mast cells, and dendritic cells. CD16 corresponds to the low-affinity Fc receptor III (FcγRIII), while CD32 corresponds to Fc receptor II (FcγRII). These receptors bind antibody-antigen immune complexes, linking innate and adaptive immunity and mediating adaptive immune responses. In research, antibodies against CD16/CD32 are commonly used to block Fc receptor-mediated interactions, preventing non-specific binding of antibodies or immunoglobulin complexes to immune cells during experiments such as flow cytometry and immunohistochemistry, thereby improving experimental accuracy.

PRODUCT DATA



BALB/c Mouse splenocytes were stained with iF700 Anti-Mouse CD16/32 clone 2.4G2 (color-filled histogram).

This product is supplied subject to the terms and conditions at www.innocyto.com/web/terms.php and may only be used as provided in the stated terms. Products are for Research Use Only.