www.innocyto.com

InnoCyto Inc.

15375 Barranca Pkwy, Suite I-103
Irvine, CA 92618
Email: info@innocyto.com
Phone: 1-949-418-7543

FITC Anti-Human CD1c Antibody

Catalog Number:	120107, 120108
Size:	25 tests, 100 tests
Target Name:	CD1C, M241, BDCA-1, R7
Regulatory Status:	RUO

PRODUCT DETAILS

Clone:	L161
Application:	Flow Cytometry
Reactivity:	Human
Format:	FITC
Isotype:	mouse IgG1
Antibody Type:	Monoclonal
Formulation:	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 0.2% (w/v) BSA
Protein Concentration:	Supplied at a lot-specific concentration.
Storage&Handling:	The antibody solution should be stored undiluted between 2°C and 8°C, and protected from prolonged exposure to light. Do not freeze.
Recommended Usage:	For flow cytometric staining, it is recommended to use 5 µL of this reagent per 0.5-1.0 million cells in a 100 µL volume. Optimal reagent performance should be determined by titration for each specific application. FITC has an excitation max at 493 nm and an emission max at 525 nm.
Excitation Laser:	Blue Laser (488 nm)
Isotype Controls:	301415
Antibody Family:	Human Antibodies
Storage & Handling:	The antibody solution should be stored undiluted between 2°C and 8°C, and protected from prolonged exposure to light. Do not freeze.
Isotype Control:	301415


BACKGROUND INFORMATION

Human CD1c, also known as BDCA-1 or CD1c antigen, is a member of the CD1 family of major histocompatibility complex (MHC) class I-like molecules that specializes in presenting lipid and glycolipid antigens to T cells. CD1c is expressed primarily on conventional dendritic cells (cDC2), B cells, monocytes, and some thymocytes. Unlike classical MHC molecules that present peptide antigens, CD1c binds lipid-based molecules derived from microbes or host tissues, thereby bridging innate and adaptive immunity. CD1c plays an important role in initiating immune responses against bacterial pathogens, particularly mycobacteria, and contributes to immune surveillance and the maintenance of self-tolerance.

Structurally, CD1c is a type I transmembrane glycoprotein composed of three extracellular α domains (α1, α2, and α3) that associate non-covalently with β2-microglobulin. The α1 and α2 domains form a deep hydrophobic antigen-binding groove that accommodates amphipathic lipid molecules, with their hydrophobic tails buried within the groove and polar head groups exposed for recognition by T-cell receptors. CD1c traffics through endosomal compartments where lipid loading and exchange occur before the molecule is transported to the cell surface. Known ligands include mycobacterial lipids such as phosphomycoketides and mannosyl phosphomycoketides, lipopeptides, phospholipids, sulfatides, and several endogenous self-lipids.

Altered CD1c expression and function have been associated with infectious diseases, autoimmune disorders, inflammatory conditions, and cancer. Defects in CD1c-mediated antigen presentation may impair antimicrobial immunity, whereas inappropriate presentation of self-lipids may contribute to autoimmune responses. In oncology, CD1c-positive dendritic cells are associated with enhanced antitumor immune responses and improved prognosis in several solid tumors. Therapeutically, CD1c is being investigated as a target for dendritic cell-based vaccines, lipid antigen immunotherapy, and strategies to enhance antigen presentation. In addition, monoclonal antibodies against CD1c are widely used to identify and isolate dendritic cell subsets for immunological research and the development of personalized cancer immunotherapies.

PRODUCT DATA



Human peripheral blood mononuclear cells were stained with iF647 Anti-Human CD19 clone CyteA-SJ25C1 and FITC Anti-Human CD1c clone L161 (left) or an isotype control (right).

This product is supplied subject to the terms and conditions at www.innocyto.com/web/terms.php and may only be used as provided in the stated terms. Products are for Research Use Only.