www.innocyto.com

InnoCyto Inc.

15375 Barranca Pkwy, Suite I-103
Irvine, CA 92618
Email: info@innocyto.com
Phone: 1-949-418-7543

APC Anti-Mouse CD31 Antibody

Catalog Number: 205307, 205308

Size: 25 µg, 100 µg

Regulatory Status: RUO

PRODUCT DETAILS

Clone: m031BB

Application: Flow Cytometry

Reactivity: Mouse

Format: APC

Isotype: Rabbit IgG

Antibody Type: Monoclonal

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 0.2% (w/v) BSA

Protein Concentration: 0.2 mg/mL

Storage&Handling: The antibody solution should be stored undiluted between 2°C and 8°C, and protected from prolonged exposure to light. Do not freeze.

Recommended Usage: For flow cytometric staining, it is recommended to use less than 0.25 µg of this reagent per 0.5-1.0 million cells in a 100 µL volume. Optimal reagent performance should be determined by titration for each specific application. APC has an excitation max at 650 nm and an emission max at 660 nm.

Excitation Laser: Red Laser (633 nm)

Isotype Controls: 303311

Antibody Family: Mouse Antibodies

Storage & Handling: The antibody solution should be stored undiluted between 2°C and 8°C, and protected from prolonged exposure to light. Do not freeze.

Isotype Control: 303311

BACKGROUND INFORMATION

Mouse CD31, also known as platelet endothelial cell adhesion molecule-1 (PECAM-1), is a cell-surface glycoprotein encoded by the Pecam1 gene. It belongs to the immunoglobulin superfamily and plays important roles in vascular biology, leukocyte trafficking, and immune regulation. CD31 is highly expressed on endothelial cells and is also found on platelets and several leukocyte populations, including monocytes, neutrophils, and certain lymphocyte subsets. It contributes to maintaining vascular integrity, regulating cell adhesion, and facilitating the movement of immune cells across blood vessel walls during inflammation.


CD31 is a type I transmembrane protein consisting of six extracellular immunoglobulin-like domains, a single transmembrane

segment, and a cytoplasmic tail containing signaling motifs. Its extracellular region mediates homophilic interactions between CD31 molecules on adjacent cells, particularly at endothelial cell junctions. CD31 can also participate in heterophilic interactions with other cell-surface molecules, including CD38 and integrin-associated proteins, depending on the cellular context. Its intracellular domain contains immunoreceptor tyrosine-based inhibitory motifs (ITIMs), which recruit signaling molecules such as SHP-1 and SHP-2 and regulate cellular activation.

CD31 is involved in inflammatory diseases, vascular injury, and tumor biology. During inflammation, it facilitates leukocyte transmigration across the vascular endothelium and helps regulate immune-cell responses. Altered CD31 expression or signaling has been associated with atherosclerosis, ischemia-reperfusion injury, and vascular dysfunction. In cancer, CD31 is widely used as a marker of endothelial cells and tumor angiogenesis, making it valuable for evaluating tumor blood-vessel density.

Therapeutically, CD31 is being investigated as a target for modulating leukocyte recruitment, vascular inflammation, and endothelial barrier function. Antibodies and other approaches that influence CD31 signaling may help regulate inflammatory responses and vascular injury. In cancer research, CD31-targeted staining and imaging are widely used to assess tumor angiogenesis and evaluate anti-angiogenic therapies.

PRODUCT DATA



C57BL/6 Mouse splenocytes were stained with APC Anti-Mouse CD31 clone m31BB (color-filled histogram) or an isotype control (gray histogram).

This product is supplied subject to the terms and conditions at www.innocyto.com/web/terms.php and may only be used as provided in the stated terms. Products are for Research Use Only.