www.innocyto.com

InnoCyto Inc.

15375 Barranca Pkwy, Suite I-103
Irvine, CA 92618
Email: info@innocyto.com
Phone: 1-949-418-7543

Anti-Human EGFR (Cetuximab Biosimilar)

Catalog Number:	501401, 501402, 501403
Size:	1 mg, 5 mg, 20 mg
Target Name:	EGFR, Proto-oncogene c-ErbB-1, Receptor tyrosine-protein kinase erbB-1, HER1
Regulatory Status:	RUO

PRODUCT DETAILS

Clone:	Cetuximab
Application:	Flow cytometry, animal model study
Reactivity:	Human
Format:	Liquid
Product Description:	Cetuximab Biosimilar, Human EGFR Monoclonal Antibody
Isotype:	Human IgG1
Clonality:	Recombinant
Immunogen:	Human EGFR / ErbB-1
Clone Number:	C225
Species specificity:	Human
Purity:	>95% by reducing SDS-PAGE
Grade:	In vivo
Storage Conditions:	4ºC
Maximal Shelf Life:	12 months
Synonyms:	ErbB1, HER1
Antibody Type:	Recombinant
Reactivity:	Human
GeneID:	1956

BACKGROUND INFORMATION

Cetuximab is a chimeric monoclonal antibody designed to specifically target the extracellular domain of the epidermal growth factor receptor (EGFR), also known as ErbB1 or HER1. Structurally, Cetuximab belongs to the immunoglobulin G1 (IgG1) subclass and is composed of two identical heavy chains and two identical light chains connected by disulfide bonds, creating a Y-shaped molecule with a molecular weight of approximately 152 kilodaltons (kDa). The molecule combines murine variable regions responsible for antigen recognition with human constant regions, resulting in a chimeric structure that preserves specificity while improving structural stability and compatibility with human immune components. It is produced using recombinant DNA technology

in mammalian expression systems, typically in murine myeloma cells.

The variable regions of the heavy (VH) and light (VL) chains contain complementarity-determining regions (CDRs) that define the antibody's epitope specificity. These CDRs bind with high affinity to the extracellular ligand-binding domain of EGFR, preventing the natural ligands, epidermal growth factor (EGF) and transforming growth factor-alpha (TGF-α), from interacting with the receptor. This blockade inhibits receptor dimerization and autophosphorylation at key tyrosine residues within the intracellular kinase domain, thereby disrupting downstream signaling cascades such as RAS-RAF-MAPK and PI3K-AKT pathways. These pathways are critical for the regulation of cellular proliferation, differentiation, and survival, making Cetuximab a valuable molecular tool for investigating growth factor signaling networks in experimental systems.

The Fc region of Cetuximab, derived from the human IgG1 isotype, contributes to its biophysical stability and extended serum half-life through interaction with neonatal Fc receptors (FcRn). It also retains the potential for effector functions via binding to Fc gamma receptors (FcγRs), enabling engagement with immune effector cells in research contexts.

This product is supplied subject to the terms and conditions at www.innocyto.com/web/terms.php and may only be used as provided in the stated terms. Products are for Research Use Only.